

# Correlation of Sea Surface Temperature and Bottom Water Temperature with the presence of Sperm Whales (*Physeter macrocephalus*) and Cuvier's beaked whales (*Ziphius cavirostris*) in the NE Aegean Sea, Greece

✉ bea.tintore@gmail.com

Beatriz Tintore<sup>1,2</sup>, Phoebe Moss<sup>1</sup>, Theodoros Tsimpidis<sup>1</sup>, Anastasia Miliou<sup>1</sup>, Dr. Gera Troisi<sup>2</sup>

<sup>1</sup> Archipelagos Institute of Marine Conservation, Samos, Greece

<sup>2</sup> Brunel University London, UK

## Introduction

The Mediterranean populations of *Physeter macrocephalus* (Pm) and *Ziphius cavirostris* (Zc) are facing increasing anthropogenic threats and yet little is known about their populations [1]. In comparison to other regions within the Mediterranean, the distribution and habitat preferences of these species in the NE Aegean Sea are extremely understudied [2]. Improving site specific knowledge on the distribution, abundance and habitat preferences of these cetacean species is essential for mitigating the impacting factors in this highly exploited region.


## Aims

- Explore the **relationship** between **depth, SST, SBT** and the **distribution of the species**
- Highlight the importance of **satellite data** when **modelling cetaceans**

## Methods

Presence/absence data for Pm and Zc were collected during **290 standardised boat-based surveys** along the **Ikarian trench** between **January 2017** and **August 2022**.

Fig 1. Study area and sightings of Pm (blue) and Zc (purple)



Satellite variables [3] included in the **GAM** using the **mgcv package** [4]:

- SST and SBT (**Copernicus Portal**)
- Depth (**EMODnet Portal**)

## Results

**2109 points of data** were used for the model, 138 with presence of Pm and 882 with presence of Zc. The values of SST and SBT ranged from **20.03°C - 21.89°C** for SST and from **13.61°C - 18.05°C** to SBT. The depth where these cetaceans were found ranged from **800 - 1200 meters**. Presence of both species analysed in this study were strongly influenced by **depth (p < 0.001)**. The model suggested an influence by **SST (p < 0.001)** for both species and **SBT (p < 0.1)** only for sperm whales.




Fig 2. Predicted smooth splines of the response variable presence/absence of *P. macrocephalus* and *Z. cavirostris* in function of explanatory variables

## Discussion

Exploration of the data showed that the **range of SST and SBT** was **too weak** to be relevant within the model **without taking seasonality into consideration** [5]. The results obtained from this study were inconclusive, however they indicate a **link between the presence of these species and SST, SBT and depth**. This association is likely driven by the **higher prey densities** present in colder, deeper waters [3,6]. Further, extensive research is recommended with the inclusion of additional variables such as **slope** and **seasonality** [5].



1. Rogan *et al.* Deep Sea Res. Part II Top. Stud. Oceanogr. 2017, 141

2. Cañadas *et al.* Ecol. Ind. 2018, 85

3. Martino *et al.* Ecog. 2021, 44,10

4. Dalla Rosa *et al.* Cont. Shelf Res. 2012, 36

5. Camrin *et al.* Anal. Rev. Mar. Sci. 2022, 14, 1

6. Virgili *et al.* PLoS ONE. 2021, 16, 8